

Klaus-Peter Schröder

Departamento de Astronomía de la
Universidad de Guanajuato, Mexico



Monitoring solar chromospheric emission with TIGRE: are we entering a Dalton Minimum?

monitoring solar and stellar Ca II
chromospheric emission in Guanajuato:
the 1.2m robotic telescope TIGRE

Liege, 15.6. 2018

a first in 2008/9:
an entirely inactive Sun




co-authors:
Marco Mittag
Jürgen Schmitt
.Dennis Jack.



UV from solar faculae coincides with Ca II K emission

- *integral solar irradiation and visual flux change only by 0.1%, no reliable data*
- *but the output of ultraviolet light ($\lambda=320-200\text{nm}$) is dominated by active regions and changes by several % (and more)*

A good proxy is CaII K emission, forms at about same T_e !



Impact of the solar far-UV light ($\lambda=320-200$ nm)

=> more activity = more far-UV = more stratospheric heating

=> less activity = cooler stratosphere = slower jetstream

=> Wider oscillation, less strength of jetstream and NAO, in winter cold high pressure areas build up, blocking situation

Sunspot number:


Above/below average

=>

**Winters:
moderate
extreme**


from:

**London Met-Office
Scaiffe et al. 2011**



How does a winter blocking situation work?

Despite stratosphere being cooler by only 1° at low solar activity, weaker and wider oscillating jetstream cannot move cold bubbles as easily as strong jetstream in high solar activity




Ca II K chromospheric line emission

as a proxy to chromospheric UV-fluxes:

problem: tiny !!

*advantage: ground-based observ. since decades,
photospheric model provides
accurate scale for surface flux*



The Mt. Wilson S-index to measure the CaII line emission:

$$S = \text{const.} (F_H + F_K) / (F_R + F_V)$$


1 Angstr. wide line cores H&K / 20 Angstr. wide quasi-continua, as such S is independent of transparency. Calibration by standard stars.

*Hence, S is of the order of the line core intensity over cont. intensity
Modern spectra: const. ca. 19, star-calibrations needed.*



*Advantage: S is independent of sky quality and calibration lamps,
best detection of even the smallest emission in the CaII core.*

Disadvantage: S does not directly compare with modern line fluxes!


Wider context: we continue O.C. Wilson's work

- *monitoring the Ca II K chromospheric emission variability, by „S-index“ = a measure relative to pseudo-continuum*
- *sample: over 100 stars brighter than 7 mag, spectral type F-K, plus about 40 cool giants of different activity degrees*
- *includes „the Sun as a star“ via moonlight spectra !*


From: Baliunas et al. 1995




S-index of Mt. Wilson project stars & the Sun



In 2008/9, the Sun reached the basal flux of „dead“ stars !



But: then came the unusual minimum of 2008/09, TIGRE, and the PhD of Marco....!



S-index of solar-type Maunder-Minimum stars

solar S-index in normal minimum

...THIS is, how the Sun looked like in Maunder Minimum!!




the basal-flux Sun:

*NO active regions at all!
=> any basal heating
is not from activity!
=> mechanical / dissip.
of acoustic waves ?*


*And where does all the
magnetic fine-structure
come from ?!!*

*~2% of convect. energy
is converted into random
fluxtubes (by a „local
dynamo“, see Vögler
& Schüssler 2007) and
into the minimal
X-ray flux (Schmitt '97)*

The historic work of O.C. Wilson et al., Mt. Wilson /CA



Chromospheric emission of cycle 24 is only <30% of previous maxima



Other activity indicators and what they mean:

Re (sunspot number, since nearly 200 yrs):

strong magnetic field through the **photosphere**

F10.7 (radioflux at $\lambda=10.7\text{cm}$, since ~50 years):


magnetic field volume in the **lower corona**

S-value (MWO-calibration, since nearly 50 years):

heating (magn. & mech.) of the **chromosphere**


related: variation of the far-UV flux (SOLSTICE, since 13 years), responsible for stratospheric heating by photodissipation of molecules

*Now all eyes on cycle 24: what is going on?!
Sunspot numbers R and $F10.7\text{cm}$ coronal radioflux, both
show only about 50% strength of previous cycles*




$\langle F10.7 \rangle - 65 \sim 0.7 \langle R_e \rangle$; 65 = minimal F10.7 value on entirely inactive days.
Maximum cycle 24 compares with activity in 2003 (mid-decline of cycle 23)


A 30% of previous cycle maxima strength, as of chromospheric emission (S-values) compares to the declined activity of 2005 (not of early 2003 !)



S_{basal} = 0.150, S_{max_av} = 0.193: S does not scale ! Goes deeper now



Latest: chromospheric emission is still 2-3 years away from 2008/9 min.



Conclusions:

- I) Solar activity cycle 24 appears half as strong as the past 3 cycles when seen by sunspot numbers and F10.7cm*
 - II) Chromospheric emission is EVEN lower than that maximum of cycle 24 at 30% of previous maxima, here is also a lowering of the baselevel (in minimum)*
 - III) The same effect is seen in the far-UV (200-280nm)!*
 - IV) This may be typical of a grand minimum, in which the Sun seems to enter again.*
 - V) The long-term reduced far-UV irradiation (by 1-2%) in a Grandminimum can explain northern hemisphere climate effects such as more cold winters*
- => monitor coming minimum and true solar analogues with TIGRE*


Much work remains to be done.....

Gracias!





SOLSTICE far-UV flux, chromospheric emission dominates:

today, same level as of 2004 is regained (as before)




The present cycle 24 seen in historic context:

S-Sun from Mt. Wilson and Lick



24
today: 0.17
(TIGRE)




present level as of 2004 or 1987 minimum!


S-Lick-Obs.

basal Ca II K flux level in S
(reached on some days in 2008/9)


***Inactive Mt. Wilson MS-stars ($S < 0.17$, near basal) over $Z=0.02$ evolution tracks, now adjusted for metallicity-differences:
All these stars are over 50% MS-lifetime (- -), most over 75%!
Note: NO evolved/inactive stars $< 1 M_{\text{sun}}$, age-limited.....***




*Highly active Mt. Wilson MS-stars ($S > 0.25 \dots 0.5$),
Z-adjusted, over $Z=0.02$ evolution tracks on MS:
Very young, scattered around the ZAMS (no surprise)*



*Moderate, **cyclic** Mt. Wilson MS-stars ($0.17 < S < 0.25$),
Z-adjusted, over $Z=0.02$ evolution tracks on MS:
Surprise: mostly less massive than the Sun!! ($\sim 50\%$ MS-lifetime)*



***Moderate, irregular Mt. Wilson MS-stars ($0.17 < S < 0.25$),
Z-adjusted, over $Z=0.02$ evolution tracks on MS:
Evolved between 50% and 75% of their MS-lifetime***



Comparison with theory of magnetic braking:

Reiners & Mohanty (2012, ApJ 746) find a relative intrinsic braking efficiency for the angular momentum of MS-stars of


$$dJ/J \sim R^{16/3} M^{-2/3}$$

Since on the MS (solar-type stars) we find $R \sim M^{0.7}$, and the decay-time $\tau \sim (dJ/J)^{-1}$, this yields

$$\tau \sim M^{-3} \sim \tau_{MS} !$$


Conclusion: The solar activity cycle is at its stability limit by BOTH, advanced evolution and by the mass of the Sun (empir. upper limit)

*Evolution tracks for $Z=0.02$ (left set) and $Z=0.01$ (right):
Metallicity **does matter for HRD position on the MS!**
*Holmberg et al. 2009 & Geneva-Copenhagen ubvy photom.:
Mt. Wilson stars occupy a range of $Z \sim 0.005 \dots 0.04$!**



The stellar perspective of solar activity:

Inactive Mt. Wilson MS-stars ($S < 0.17$) over $Z=0.02$ evolution tracks: more evolved than the Sun (circle)!



S-index of Mt. Wilson project stars & the Sun

